

Search for Rare Decays
 $B_s \rightarrow \mu^+ \mu^-$ and $B_d \rightarrow \mu^+ \mu^-$
Using CDF Run2 Data

Vyacheslav Krutelyov

Texas A&M University

With D.Glenzinski, M.Herndon, C.J.Lin,

T.Kamon, J.Thom, A.Yagil

For CDF Collaboration

May 28, 2004

Chicago Flavor Seminar

Fermilab

Outlines

☞ Introduction

- Theoretical motivations.
- Existing limits

☞ CDF Run2

☞ $\text{Br}(B_{s(d)} \rightarrow \mu^+ \mu^-)$ measurement.

- Blind analysis.
- Signal & Background
 - ◆ **Discriminating variables**
- Background estimation method.

☞ Acceptance and efficiency

☞ Optimization \rightarrow Expected limit

☞ Observed limit

☞ Summary and plans



Theoretical motivations. Current limits.

Flavor Changing Neutral Current.

Loop contribution only in SM.

$$\text{Br}_{\text{SM}}(B_s \rightarrow \mu^+ \mu^-) = (3.4 \pm 0.5) \times 10^{-9}$$

$$\text{Br}_{\text{SM}}(B_d \rightarrow \mu^+ \mu^-) = (1.00 \pm 0.14) \times 10^{-10} \text{ (hep-ph/0303060)}$$

Only upper experimental limit exists:

$$\text{Br}_{\text{exp}}(B_s \rightarrow \mu^+ \mu^-) < 2.0 \times 10^{-6} \text{ 90\% C.L. CDF RunI @100/pb.}$$

$$\text{Br}_{\text{exp}}(B_d \rightarrow \mu^+ \mu^-) < 1.6 \times 10^{-7} \text{ 90\% C.L. Belle '03 @78/fb.}$$



Limits are far away from the SM value: can test for a possible new physics.

MSSM: $\text{Br}(B \rightarrow \mu^+ \mu^-)$ enhanced by $\tan\beta > 10$ terms $\sim \tan^6\beta$.
Up to 100 over the SM prediction.


R-parity violating models give tree level contributions.
Not heavily constrained by other observables.

Can be seen in Run2


Other models enhance less. E.g., universal extra dimensions.
Up to +70% for $B_s \rightarrow \mu^+ \mu^-$



mSUGRA, SO(10)



CDF Run2.



- ☞ $\sqrt{s}=1.96\text{TeV}$ $\sigma(\text{Inelastic})\sim 60\text{mb}$. $L \sim 6 \cdot 10^{31}\text{cm}^{-2}\text{s}^{-1}$ ($\sim 230\text{pb}^{-1}$ good recorded)
 - Plan: $\sim 8 \cdot 10^{31}\text{cm}^{-2}\text{s}^{-1}$ (Run2a); $\sim 2 \cdot 10^{32}\text{cm}^{-2}\text{s}^{-1}$ (4-8/fb Run 2)
- ☞ 1.7MHz collision \rightarrow 20kHz L1 trigger \rightarrow 350Hz L2 \rightarrow 60Hz L3/logging rate.
 - $\sim 80\%$ L1 ($\sim 1/3$ at L3) of the triggers are B physics

Run2 vs Run1

- ☞ Better silicon coverage (x2), muon detection, improved tracking.
- ☞ Better triggers: lower track p_T , higher efficiency.

Br(B_{s(d)} → μ+μ-) measurement.

$$BR(B_s \rightarrow \mu\mu) \leq \frac{N_{CL}(n_{obs}, n_{bg})}{2 \cdot a \cdot e_{total} \cdot s_{Bs} \cdot \int L dt}$$

3% (CMU&CMP Run1)

25% Run1

CL=90% upper limit on $\langle n_{sig} \rangle$ for n_{obs} and n_{bg}

171pb-1 or 10 trillion collisions

only Run1 $\sigma_{Bs} = 0.9 \mu b$ for $p_T^{Bs} > 6 \text{ GeV}/c$, $|y| < 1$
(use this as a baseline selection)






Dimuon trigger and $p_T > 6 \text{ GeV}$
baseline sample: 2940 events

- ☞ Expect to detect at most few events that might only look like $B_{s(d)} \rightarrow \mu + \mu -$
- ☞ SM predicts 0 events \Rightarrow really a “search”
 - emphasis on performing an unbiased optimization
- ☞ Don't look at the data signal region (blind search)
 - Signal inside $5.169 < M_{\mu\mu} < 5.469 \text{ GeV}/c^2$ ($\pm 3\sigma$ window)
- ☞ demonstrate understanding of background events
- ☞ accurately estimate α (acceptance) and ϵ (efficiency)
- ☞ intelligently optimize cuts



Signal & Background

Need to select discriminating variables that leave most of the signal and remove most of the background.



Most Promising Discriminating Variables

- Invariant mass
- $c\tau$: $L_{xy} \frac{M}{P_t(B)}$ ← 2D lifetime $\times c$
- $\Delta\Phi$: $\phi(\vec{p}_T^B) - \phi(vtx)$
- Isolation $\frac{P_t(B)}{P_t(B) + \sum_{\Delta R < 1} P_t(trk)}$

Largest expected background:
gluon splitting $g \rightarrow b\bar{b} \rightarrow \mu\mu$

Background estimation (1)

$$N(\text{bgd}) = N(\text{sideband} | c\tau, \Delta\Phi) * R(\text{iso}) * R(M_{\mu\mu})$$

- $N(\text{sideband} | c\tau, \Delta\Phi)$ == sideband events passing $c\tau$ and $\Delta\Phi$ cuts
- $R(\text{iso})$ == fraction of bgd expected to survive an isolation cut
- $R(M_{\mu\mu})$ == given $N(\text{sideband})$, estimate of $N(\text{signal window})$

$$r_{xy} = \frac{1}{N-1} \cdot \frac{\sum (x_i - \hat{x})(y_i - \hat{y})}{s_x s_y}$$

⇒ Only $c\tau$ & $\Delta\Phi$ are significantly correlated ($\rho \sim -0.3$),


the rest is not ($|\rho| \sim 0.05$) ⇒ can

determine $R(\text{iso})$ and $R(M_{\mu\mu})$

on sample w/ NO $c\tau$, $\Delta\Phi$ cuts

⇒ $M_{\mu\mu}$ is linear ⇒ can use

sidebands to estimate background



Background estimation (2)

⇒ Method cross-checks on control samples

Cuts	N(exp)	N(obs)	$P(>=obs exp)$
loose	23 ± 3	27	20%
optimal	8 ± 1	11	17%
tight	1.2 ± 0.3	2	34%

- Sample = outside signal window
- $P(>=obs|exp)$: Poisson probability of observing $\geq obs$ events with exp expected

⇒ $B \rightarrow hh$ background is negligible.

Will be sensitive to these when the limit reaches $O(10^{-9})$

Acceptance and efficiency

$$\mathbf{a} \cdot \mathbf{e}_{\text{total}} = \mathbf{a} \cdot \mathbf{e}_{\text{trig}} \cdot \mathbf{e}_{\text{reco}} \cdot \mathbf{e}_{\text{zvtx}} \cdot \mathbf{e}_{\text{disVars}}$$

☞ α – acceptance = $6.6 \pm 0.5\%$

➤ so far include only muons with $|\eta| < 0.6$ (CMU&CMP)

☞ $e_{\text{total}} \approx \underline{0.52} \cdot e_{\text{disVars}}$

➤ take trigger and reconstruction efficiency from J/ψ data

◆ +/- 10% (relative) syst due to kinematic difference with $B_{s(d)}$

➤ $e_{\text{trig}} = 85 \pm 3\%$

➤ $e_{\text{reco}} = (e_{\text{COT}} e_{\mu\text{reco}} e_{\mu\text{qual}} e_{\text{SVX}})^2 = 71 \pm 3\%$

➤ $e_{\text{zvtx}} = 95.1 \pm 0.1 \pm 0.5\%$ -- luminous region selection

☞ e_{disVars} – efficiency of discriminating variables cuts, changes during optimization:

➤ varied within 28-78% for ~200 combinations of $(M_{\mu\mu}, c\tau, \Delta\Phi, \text{Iso})$.

➤ cross-check the method using $B^+ \rightarrow J/\psi K^+$ data and MC ($\pm 5\%$ relative syst)

Optimization Results

Optimize on $(M_{\mu\mu}, c\tau, \Delta\Phi, \text{Iso})$ to get the best expected limit

$$\langle \text{BR}(B_s \rightarrow \mu\mu) \rangle = \frac{\sum_n N(n | n_{\text{bg}}) \cdot P(n | n_{\text{bg}})}{2 \cdot a \cdot e_{\text{total}} \cdot s_{B_s} \cdot \int L dt}$$

Poisson prob

$(c\tau, \Delta\Phi, \text{Iso}) =$

$(>200 \text{ mm}, <0.10 \text{ rad}, >0.65)$

and mass window $\pm 80 \text{ MeV}$ around

world avg $B_s(d)$: 5.369 GeV (5.279 GeV)


$\triangleright B_s(d)$: $\alpha \times \epsilon_{\text{total}} = (2.0 \pm 0.2)\%$

$(\alpha \approx 6.6\%, \epsilon_{\text{total}} \approx 30\%)$

\triangleright Accepted bgd $\sigma = (6 \pm 2) \text{ fb}$

- Expected background

$\langle \text{Bgd} \rangle$ in $171 \text{ pb}^{-1} = 1.1 \pm 0.3 \text{ events}$



Observed limits


$BR(B_s \rightarrow \mu^+ \mu^-) < 5.8 \times 10^{-7}$ 90% C.L.
(7.5×10^{-7} 95% C.L.)

3 times better than Run I (previous world best)

$BR(B_d \rightarrow \mu^+ \mu^-) < 1.5 \times 10^{-7}$ 90% C.L.
(1.9×10^{-7} 95% C.L.)

slightly better than (1.6×10^{-7} 90% C.L., just published)

B-factory/Belle result





Conclusions for SuperSymmetry (@95% C.L.)

SO(10)

R. Dermisek *et al.*,
hep-ph/0304101

R-parity violating

R. Arnowitt *et al.*,
PLB 538 (2002) 121,
new plot by B.Dutta




New limit excludes **significant** amounts of the previously allowed SUSY models parameters

Summary

- ☞ CDF collaboration produces competitive measurements of rare B dimuon decays.
- ☞ $\text{Br}(B_s \rightarrow \mu\mu) < 5.8 \times 10^{-7}$ at 90% C.L.
- ☞ $\text{Br}(B_d \rightarrow \mu\mu) < 1.5 \times 10^{-7}$ at 90% C.L.
- ☞ This result is submitted to (and accepted by) PRL
 - [hep-ex/0403032](https://arxiv.org/abs/hep-ex/0403032)
- ☞ We expect to have $\text{Br}(B_{d(s)} \rightarrow \mu\mu)$ measurement significantly improving this result later this year.
 - This result does not include muons from CMX muon subdetector: another 6% in acceptance (doubling the present acceptance)
 - ◆ has larger backgrounds
 - More integrated luminosity is being collected (already have +50%)
 - Have a slightly improved reconstruction software

(c) Copyright Tony Frampton 2001

DR. DIDD



AND ON THAT EXCITING NOTE...



BACKUP SLIDES

Data Samples

- $OS+$: opposite-sign muon pairs, $c\tau > 0$
our signal sample - not used for xchecks
- $OS-$: opposite-sign muon pairs, $c\tau < 0$
- $SS+$: same-sign muon pairs, $c\tau > 0$
- $SS-$: same-sign muon pairs, $c\tau < 0$


Motivations: $B_s \rightarrow \mu^+ \mu^-$ (mSUGRA)

R. Arnowitt *et al.*,
 hep-ph/0203069,
 PLB 538 (2002) 121

Overlap with measured da_μ (BNL) in mSUGRA parameter space.

Overlap with dark matter=LSP allowed region.




Eliminate large parameter space (~ all for $\tan\beta > 40$), with $Br(B_s \rightarrow \mu^+ \mu^-) \sim 10^{-8}$ in Run2 (15/fb)



R_p Violation: Br vs. $m_{1/2}$


R. Arnowitt *et al.*,
 hep-ph/0203069,
 PLB 538 (2002) 121

e.g., $W_{TRPV} = l_{ijk} L_i L_j E_k + l'_{ijk} L_i Q_j D_k + l''_{ijk} U_i D_j D_k$



B → hh PR plot

now ~1K events with
180/pb



$B_{d(s)} \rightarrow mm$: Run I Results

CDF, PRD 57, 3811 (1998)

$\int L dt = 98 \text{ pb}^{-1}$ ($\sim 5 \times 10^{12}$ collisions)

B_d : 5.205-5.355 GeV/c^2

B_s : 5.300-5.450 GeV/c^2

$N_{\text{obs}} = 1$ with 5.344 GeV/c^2

(consistent with an expectation from the Standard Model backgrounds)

$Br(B_d \rightarrow mm) < 8.6 \times 10^{-7}$ (95% C.L.)

$Br(B_s \rightarrow mm) < 2.6 \times 10^{-6}$ (95% C.L.)

